

AD-A167 407

THE EFFECTS OF CYCLOPEGIA ON THE VISUAL CONTRAST  
SENSITIVITY FUNCTION(U) ARMY AEROMEDICAL RESEARCH LAB  
FORT RUCKER AL W G BACHMAN ET AL. FEB 86 USARL-86-2


1/1

UNCLASSIFIED

F/G 6/5

NL





MICROCOPY

CHART

12



USAARL Report No. 86-2

AD-A167 407

**THE EFFECTS OF CYCLOPEGIA ON THE VISUAL  
CONTRAST SENSITIVITY FUNCTION**

By  
**William G. Bachman  
Isaac Behar**

**DTIC**  
**ELECTE**  
**MAY 05 1986**  
**S** **D**

**SENSORY RESEARCH DIVISION**

**February 1986**

DTIC FILE COPY

86 5 5 020

Approved for public release, distribution unlimited

**USAARL**

## NOTICE

### Qualified Requesters

Qualified requesters may obtain copies from the Defense Technical Information Center (DTIC), Cameron Station, Alexandria, Virginia 22314. Orders will be expedited if placed through the librarian or other person designated to request documents from DTIC.

### Change of Address

Organizations receiving reports from the US Army Aeromedical Research Laboratory on automatic mailing lists should confirm correct address when corresponding about laboratory reports.


### Disposition

Destroy this report when it is no longer needed. Do not return it to the originator.

### Disclaimer

The views, opinions, and/or findings contained in this report are those of the author and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other official documentation. Citation of trade names in this report does not constitute an official Department of the Army endorsement or approval of the use of such commercial items.

Reviewed:



BRUCE C. LEIBRECHT  
LTC, MS  
Director, Sensory Research  
Division

Released for Publication:



J. D. LaMOTHE  
LTC, MS  
Chairman, Scientific  
Review Committee



DUDLEY R. PRICE  
Colonel, MC  
Commanding

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER USAARL Report No. 86-2	2. GOVT ACCESSION NO. <b>AD A167402</b>	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle)  The Effect of Cycloplegia on the Visual Contrast Sensitivity Function		5. TYPE OF REPORT & PERIOD COVERED
7. AUTHOR(s)  William G. Bachman and Isaac Behar		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Sensory Research Division US Army Aeromedical Research Laboratory Fort Rucker, AL 36362-5000		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS US Army Medical Research and Development Command Fort Detrick Frederick, MD 21701-5012		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS  3E162777A879,164
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE February 1986
		13. NUMBER OF PAGES 22
		15. SECURITY CLASS. (of this report)  UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  Contrast Sensitivity Function Cycloplegia Glare Vision		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  See reverse side.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

20. ABSTRACT.

Contrast sensitivity assessment is one of several emergent techniques being considered for inclusion in a visual standards test battery for the Army, particularly for the evaluation of Army aviators. Since a cycloplegic refraction is required for initial selection of candidates for Class I and Class IA flying duty, it is important to determine what effect, if any, cycloplegia has on the contrast sensitivity function. Twelve subjects, officers in preparation for flight training, who had passed a recent Class I flight physical, were tested. Contrast sensitivity functions were obtained under normal ambient conditions and in the presence of a glare source, both under manifest and cycloplegic conditions. Cycloplegia produced a small reduction in contrast sensitivity under normal ambient conditions, and a greater reduction under glare conditions. For both conditions, the cycloplegia effect was greater for the higher spatial frequency gratings than for the lower.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

### Acknowledgments

The authors wish to acknowledge the valued contributions to this project of LTC Bruce Leibrecht, Ms. Carolyn Johnson, SGT Rosalinda Ibanez, SSG Nonilon Fallaria, and SGT Marshall Smith.

Accession For	
NTIS CRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	



## TABLE OF CONTENTS

	PAGE NO.
Introduction .....	3
Method .....	5
Results .....	8
Discussion .....	11
Conclusions .....	11
References .....	12
Appendixes	
A. Repeated measures ANOVA summary table .....	15
B. List of manufacturers .....	16



## LIST OF TABLES

TABLE NO.		PAGE NO.
I	Mean optical correction .....	5
II	Mean pupillary diameter .....	5

## LIST OF FIGURES

FIGURE NO.		PAGE NO.
1	Contrast sensitivity functions for manifest and cycloplegic conditions directly com- pared .....	8
2	Contrast sensitivity functions for glare and no glare conditions directly compared .....	10

## INTRODUCTION

Contrast sensitivity assessment has become a major tool for evaluating human spatial vision (Schade, 1956; Campbell, 1983). The contrast sensitivity function (CSF) characterizes the threshold sensitivity of the visual system to sinusoidal variations in contrast over a wide range of target sizes or spatial frequencies. The performance of the visual system is measured over an entire range rather than at specific high frequencies as is done in Snellen letter or Landolt C visual acuity determination.

Comerford (1983) stated that the CSF provides several advantages over other vision tests commonly used in clinical practice. First, it provides a measure of the integrity of both central and peripheral vision requiring similar judgments on the part of the patient. Second, for low spatial frequency stimuli, the detection of the grating requires the integration of visual information over a large expanse of the retina. Third, while better acuity generally indicates the patient's ability to see small details in the environment, the CSF also measures the ability to see large details. These details, which are analogous to low frequency sensitivity, provide input to facial recognition, figure-ground judgments, and other important information relating to the ability to function in the environment. Fourth, the CSF is a sensitive indicator of small differences in visual function.

The ability to detect small differences in visual function can have a significant application in the military, especially in the aviation environment. For example, Ginsburg et al. (1982) and Ginsburg, Easterly, and Evans (1983) demonstrated that contrast sensitivity was found to be better than visual acuity for predicting a pilot's performance in detecting small low contrast targets both in aircraft simulators and in the field. Contrast sensitivity testing also is being conducted by NASA on space shuttle flights to determine the effects of low gravity on vision.

Contrast sensitivity testing currently is used in the Army primarily in the area of research. It is used in special medical cases to determine the effect on visual function of eye pathology, e.g., an aviator with a developing cataract. Contrast sensitivity assessment is one of several emergent techniques being considered for inclusion in a visual standards test battery for the Army. This battery would offer the potential for characterizing individual differences not now captured with standard visual testing.

Instruments and methods for contrast sensitivity determination have existed almost exclusively in the research environment. Arden (1978) developed the first practical clinical test of contrast sensitivity using gratings printed on plates. A clinical contrast sensitivity screening instrument has been introduced (Ginsburg et al., 1984), as well as an automated refractor which measures contrast sensitivity and displays results digitally as visual acuity (CooperVision Dicon AR5000\*). Also now available to the clinician are spatial frequency charts similar to acuity charts which are designed to be used for both distance and near testing (Ginsburg, 1984). Therefore, contrast sensitivity testing is becoming more useful in clinical vision evaluation and screening.

Because of growing acceptance and ease of use, as well as practical importance to Army aviation, contrast sensitivity testing would appear to be an important tool in the evaluation of potential Army aviators. Army Regulation 40-501, Change 34, indicates that a cycloplegic refraction is required for initial selection of candidates for Class I and Class IA flying duty. Cycloplegia, in this case, is defined by Cline, Hoffstetter, and Griffin (1980) as an artificially induced paralysis of the ciliary muscle and the power of accommodation, usually accompanied by a dilated pupil. It is relevant therefore to determine what effect, if any, cycloplegia has on the contrast sensitivity function.

Several previous studies have examined the effects on the CSF by drugs that modify the pupillary or accommodative dynamics. Singh et al. (1981) determined that mydriasis alone without paralysis of accommodation did not affect contrast sensitivity in normal older (50 to 84 years) observers. Baker et al. (1983) found that for subjects who had been given atropine intramuscularly, there was a small loss of sensitivity at the highest tested spatial frequency (20 cycles per degree). Campbell and Green (1965) demonstrated the effect on contrast sensitivity of various artificial pupil sizes under atropinized conditions. Using neutral density filters to compensate for the change in retinal illumination due to the changes in pupil area, they found a progressive reduction of contrast sensitivity as the pupil was made larger. Kay and Morrison (1985) replicated the Campbell and Green study except that changes in retinal illumination associated with changes in pupil size were not compensated for in order to simulate natural viewing. They found that the low spatial frequency portion of the CSF "was only

-----  
\* See Appendix B

marginally affected by pupil diameter" as well as the high spatial frequency portion which "was also relatively unaffected by pupil diameter."

The contrast sensitivity function measured under normal clinical conditions does not provide information for all situations encountered by a subject in his or her daily environment. Paulsson and Sjostrand (1980) used a bright light glare source introduced into the visual field in order to enhance the effects of intraocular light scattering on the CSF. In the same vein, Carney and Jacobs (1984) stated that the contrast sensitivity function can be sensitized by the presence of a glare source to allow a more accurate determination of any visual loss.

The purpose of this study then was to evaluate the effect of cycloplegia on the contrast sensitivity function. This was done in a manner simulating clinical conditions as closely as possible as well as with the introduction of a glare source. Subjects were candidates for flight training at Fort Rucker, Alabama.

## METHOD

### Subjects

Twelve subjects were used, 11 males and 1 female. All were aged 22 or 23, except one who was 27. All were officers in preparation for flight training at the Army Aviation Center at Fort Rucker, Alabama, who had passed a recent Class I flight physical, and were free of eye disease. All subjects received an intraocular pressure test (Reichert\* noncontact tonometer) and were within normal limits.

### Procedures

Subjects were refracted at a distance of three meters using both standard subjective refraction and static retinoscopy to determine spherical and cylindrical components. Refraction was accomplished under both manifest (undilated) and cycloplegic (dilated) conditions. See Table I for the mean optical correction under both conditions. The subjects then wore the best possible optical correction (spectacles) to minimize blur at 3 m for whichever condition was being tested. Optical corrections, to include plano results, were provided using a standard trial frame. All subjects resolved 20/20 or 20/15 with each eye unaided as well as with spectacle corrections. Acuties were measured using the Baylor Video Acuity Tester (B-VAT)\* which has a 12" diagonal CRT for video display of target sizes from

20/10 to 20/400. Mean luminance of the screen was held at 26.5 fL which was identical to the mean luminance of the contrast sensitivity testing screen.

Pupil diameters were measured for all subjects. Diameters were measured with a PD rule to the nearest 0.5 mm while the subject viewed the contrast sensitivity display. This was accomplished with and without glare under both undilated and dilated conditions. See Table II for the mean pupillary diameters.

Diopters		
	Sphere	Cylinder
Manifest	+0.09	-0.36
Cyclopegic	+0.46	-0.34

TABLE I. Mean optical correction.

	Manifest	Cyclopegic
No glare	4.92	8.08
Glare	3.08	8.08

TABLE II. Mean pupillary diameter.

Cycloplegia was induced using 1 percent Cyclogel\* (cyclopentolate) which is a diagnostic parasympatholytic drug administered directly in the eye. Each subject received one drop in each eye followed by a second drop after 5 minutes. Cyclopentolate blocks the responses of the sphincter muscle of the iris and the accommodative muscle of the ciliary body to cholinergic stimulation, producing pupillary dilation (mydriasis) and paralysis of accommodation (cycloplegia).

#### Contrast Sensitivity Measurement

Testing was conducted in a room in which all surfaces, walls, ceiling, and floor were matte black. Room illumination was provided by four recessed ceiling incandescent lamps adjusted to provide 12 fc at the observer's table. The contrast sensitivity functions were obtained with a Nicolet Optronics CS2000 Contrast Sensitivity Testing System\*. The video display had a mean luminance of 26.5 fL, and at the 3-meter viewing distance, subtended 4.4 degrees by 5.6 degrees. For a glare source, the display was surrounded by a high intensity (4300 fL) fluorescent lamp (Aristo DA-17\*) which was masked so that no direct light reached the screen. The choice of a surrounding glare source instead of a more commonly used laterally placed small glare source was based on the findings of Miller et al. (1972) that the former was less fatiguing and helped the subjects maintain fixation on the centrally located display.

The contrast threshold was measured using a variation of the method of increasing contrast (Ginsburg and Cannon, 1983), which is similar to an ascending method of limits psychophysical procedure. On each trial, the display contrast began near zero and after a variable delay increased under computer control uniformly at a rate at which 50 percent contrast would be reached in 45 s. The subject's task was to depress a response switch immediately upon detecting the emergence of a grating pattern on the display screen. Each subject was tested on 3 separate days, the first of which was for training. For half of the subjects, testing on the second and third day was under manifest and cycloplegic conditions, respectively, while for the remaining subjects the order of conditions was reversed. On the training day, they received verbal instructions followed by 18 practice trials consisting of three trials at each spatial frequency: 0.5, 1, 2, 4, 8, and 16 cpd, in an intermixed random order. This then was repeated with the glare source turned on. On each test day, the subject received five warm-up trials (one trial each at 0.75, 1.5, 3, 6, and 12 cpd in random order) followed by a random series of 36 trials consisting of 6 trials at each of the spatial frequencies 0.5 to 16 cpd. Following a short break, this 41-trial set was repeated with the glare source turned on. For each subject, mean log threshold contrast was calculated for each

combination of conditions: manifest-glare source off, cycloplegic-glare source off, manifest-glare source on, cycloplegic-glare source on.

## RESULTS

The mean contrast sensitivity (reciprocal of group mean contrast threshold) as a function of spatial frequency is presented graphically in Figure 1. The upper left panel summarizes the results for the two conditions with the glare source turned off, while the upper right panel summarizes the results with the glare source turned on. For both no glare ( $F=5.76$ ,  $df=1,11$ ,  $p<.036$ ) and glare ( $F=9.79$ ,  $df=1,11$ ,  $p<.01$ ), contrast sensitivity is superior under manifest compared to cycloplegic conditions for all spatial frequencies except the lowest spatial frequency (0.5 cpd) under glare. The log ratios of the mean contrast thresholds for each spatial frequency with




Figure 1. Contrast sensitivity functions (upper panels) and visuograms (lower panels) obtained under manifest and cycloplegic conditions, with and without glare. CSFs for manifest and cycloplegic conditions are directly compared.

and without cycloplegia constitute visuograms (Lundh and Arlinger, 1984), and are plotted in the lower panels of this figure.

In the absence of glare, the loss in contrast sensitivity caused by cycloplegia is minimal for the lower spatial frequencies, but increases monotonically between 2 and 16 cpd. In the presence of glare, the effect of cycloplegia in reducing sensitivity is somewhat greater (except for 0.5 cpd) than it was without glare, and the reduction with increasing spatial frequency is essentially monotonic. An overall repeated measures analysis of variance was performed on these data and a summary table is presented in Appendix A. The main effects of cycloplegic condition ( $F=8.92$ ,  $df=1,11$ ,  $p=.012$ ) and spatial frequency ( $F=177.17$ ,  $df=5,55$ ,  $p<.0001$ ) are highly significant, while the main effect of glare is not significant.

However the interaction of glare and cycloplegic condition, is significant ( $F=6.03$ ,  $df=1,11$ ,  $p=.032$ ), reflecting the greater separation of the manifest and cycloplegic conditions in the presence of glare rather than in its absence. A significant interaction between cycloplegic condition and spatial frequency ( $F=2.95$ ,  $df=5,55$ ,  $p=.020$ ) reflects the greater separation of the manifest and cycloplegic conditions at the higher spatial frequencies than at the lower spatial frequencies. Finally, a significant interaction between glare and spatial frequency ( $F=7.74$ ,  $df=5,55$ ,  $p<.0001$ ) reflects the superior contrast sensitivity in the presence of glare at the lowest spatial frequency (0.5 cpd), but reduced contrast sensitivity in the presence of glare at the intermediate spatial frequencies (2 to 8 cpd), replicating a pattern of results previously found (Behar, 1984).

The comparison of the effects of glare versus no glare is facilitated by the replot of the data in Figure 2. It can be seen that contrast sensitivity is superior in the presence of glare at the lowest spatial frequency (0.5 cpd) when tested under both manifest and cycloplegic conditions. For the remaining spatial frequencies, important differences exist. Under manifest conditions, with the eye in its normal physiological state, no glare sensitivity is evident; contrast sensitivity is equivalent in the presence of the glare source and in its absence. On the other hand, under cycloplegic conditions, considerable glare sensitivity is evident, especially in the middle spatial frequencies (2 to 8 cpd).






Figure 2. Contrast sensitivity functions (upper panels) and visuograms (lower panels) obtained under manifest and cycloplegic conditions, with and without glare. CSFs for glare and no glare conditions are directly compared.

## DISCUSSION

The purpose of this study was to evaluate the effect of cycloplegia on the contrast sensitivity function in a group that is required to have a cycloplegic refraction in order to pass a flight physical. The results indicate that there is a small, but significant loss of sensitivity under dilated conditions and that this loss is magnified by the introduction of glare. Since subjects wore corrections for the test viewing distance, these reductions can be attributed primarily to differences in pupil size in the various conditions. An increase in the size of the pupil increases retinal illuminance, thus would be expected to improve contrast sensitivity especially to higher spatial frequencies (Owsley et al., 1985; Wright and Drasdo, 1985). On the other hand, the larger pupil suffers greater levels of aberrations and reduced depth of field resulting in impairment of contrast sensitivity (Campbell and Gubisch, 1966). When both factors were allowed to operate in opposition, as in the present study, in that of Kay and Morrison (1985), and in that of Singh et al. (1981), relatively small changes in the CSF were found. Singh's subjects were between the ages of 50 and 84 years, so would be expected to have reduced pupils associated with senile meiosis (Said and Sawires, 1972). The mydriatic induced dilated pupil in these subjects resulted in a very large increase in retinal illumination that should favor an improvement in the CSF; however, the degradation in retinal image quality accompanying the larger pupil evidently exactly canceled the illumination advantage leaving the CSF unchanged. In the present study, since the subjects were only in their 20s, the increase in retinal illumination was considerably less, so the more potent factor was the reduction in retinal image quality, resulting in a small net impairment of the CSF.

## CONCLUSIONS

If the contrast sensitivity function is to be obtained during the flight physical, it should be obtained prior to the administration of cycloplegia. If the CSF is determined after cycloplegia, then it is most important to avoid viewing with a glare source in the field of view (such as a desk lamp or uncovered window) in order to avoid a biased assessment.

## REFERENCES

- Department of the Army. 1983. Army Regulation 40-501, Change 34. Medical services standards of medical fitness. Headquarters, Department of the Army, Washington, DC.
- Arden, G.B. 1978. The importance of measuring contrast sensitivity in cases of visual disturbance. British Journal of Ophthalmology. 62:198-209.
- Baker, R., Adams, A., Jampolsky, A., Brown, B., Haegerstrom-Portnoy, G., and Jones, R. 1983. Effects of atropine on visual performance. Military Medicine, 148:530-535.
- Behar, I. 1984. The effects of haze and glare on visual contrast sensitivity--preliminary results. In Proceedings of the Tri-Service Aeromedical Research Panel Fall Technical Meeting. NAMRL Monograph - 33, November, 1984.
- Campbell, F.W. 1983. Why do we measure contrast sensitivity? Behavioral Brain Research. 10:87-97.
- Campbell, F.W. and Green, D.G. 1965. Optical and retinal factors affecting visual resolution. Journal of Physiology. 181:576-593.
- Campbell, F.W. and Gubisch, R.W. 1966. Optical quality of the human eye. Journal of Physiology (London). 186: 558-578.
- Carney, L.G. and Jacobs, R.J. 1984. Mechanisms of visual loss in corneal edema. Archives of Ophthalmology. 102: 1068-1071.
- Cline, D., Hofstetter, H.W., and Griffin, J.R. 1980. Dictionary of Visual Science. Radnor, PA: Chilton Book Company.
- Comerford, J.P. 1983. Vision evaluation using contrast sensitivity functions. American Journal of Optometry & Physiological Optics. 60:394-398.
- Ginsburg, A.P. 1984. A new contrast sensitivity vision test chart. American Journal of Optometry & Physiological Optics. 61:403-407.

- Ginsburg, A.P., Evans, D.W., Sekuler, R., and Harp, S.A. 1982. Contrast sensitivity predicts pilots' performance in aircraft simulators. American Journal of Optometry & Physiological Optics. 59:105-109.
- Ginsburg, A.P., Easterly, J., and Evans, D.W. 1983. Contrast sensitivity target detection field performance of pilots. Proceedings, Human Factors Society.
- Ginsburg, A.P., and Cannon, M.W. 1983. Comparison of three methods for rapid determination of threshold contrast sensitivity. Investigative Ophthalmology and Visual Science. 24:798-802.
- Ginsburg, A.P., Evans, D.W., Cannon, M.W., Jr., Owsley, C., and Mulvanny, P. 1984. Large-sample norms for contrast sensitivity. American Journal of Optometry & Physiological Optics. 61:80-84.
- Kay, C.D. and Morrison, J.D. 1985. The effects of pupil size and defocus on contrast sensitivity in man. Journal of Physiology. 367,15P.
- Lundh, B.L. and Arlinger, S. 1984. Three dB-scales for the standardized visuogram. Vision Research. 24:889-890.
- Miller, D., Jernigan, M.E., Molnar, S., Wolf, E., and Newman, J. 1972. Laboratory evaluation of a clinical glare tester. Archives of Ophthalmology. 87:324-332.
- Owsley, C., Gardner, T., Sekuler, R., and Lieberman, H., 1985. Role of the crystalline lens in the spatial vision loss of the elderly. Investigative Ophthalmology and Visual Science. 26:1165-1170.
- Paulsson, L.-E. and Sjostrand, J. 1980. Contrast sensitivity in the presence of a glare light theoretical concepts and preliminary clinical studies. Investigative Ophthalmology and Visual Science. 19:401-406.
- Said, F.S. and Sawires, W.S. 1972. Age dependence of changes in pupil diameter in the dark. Optica Acta. 19: 359-361.
- Schade, O.H. 1956. Optical and photoelectric analog of the eye. Journal of the Optical Society of America. 46: 721-739.

Singh, H., Cooper, R.L., Alder, V.A., Crawford, G.J., Terrell, A., and Constable, I.J. 1981. The Arden grating acuity: effect of age and optical factors in the normal patient, with prediction of the false negative rate in screening for glaucoma. British Journal of Ophthalmology. 65:518-524.

Wright, C.E. and Drasdo, N. 1985. The influence of age on the spatial and temporal contrast sensitivity function. Documenta Ophthalmologica. 59:385-395.

REPEATED MEASURES ANOVA SUMMARY TABLE

SOURCE	SUM OF SQUARES	DEGREES OF FREEDOM	MEAN SQUARE	F	TAIL PROB.	GREENHOUSE-GEISSER PROB.	HUYNH-FELDT PROB.
1 MEAN ERROR	1079.16514 2.56705	1 11	1079.16514 0.26973	4000.89	0.0000		
2 CYCLO ERROR	0.36894 0.45490	1 11	0.36894 0.04135	8.92	0.0124		
3 GLARE ERROR	0.00451 0.21687	1 11	0.00451 0.01972	0.23	0.6317		
4 CG ERROR	0.03777 0.06893	1 11	0.03777 0.00627	6.03	0.0320		
5 SPAFREQ ERROR	25.35698 1.57436	5 55	5.07140 0.02862	177.17	0.0000	0.0000	0.0000
6 CS ERROR	0.13308 0.49622	5 55	0.02662 0.00902	2.95	0.0193	0.0452	0.0252
7 GS ERROR	0.25264 0.37224	5 55	0.05053 0.00677	7.47	0.0000	0.0013	0.0003
8 CGS ERROR	0.05753 0.30885	5 55	0.01151 0.00562	2.05	0.0959	0.1354	0.1137

## EPSILON FACTORS FOR DEGREES OF FREEDOM ADJUSTMENT

GREENHOUSE-GEISSER	HUYNH-FELDT
0.3804	0.4574
0.6175	0.8860
0.5082	0.6737
0.5263	0.7068

ERROR TERM
5
6
7
8

## APPENDIX B

Alcon Laboratories, Inc.  
P. O. Box 1959  
6201 South Freeway  
Fort Worth, TX 76134

Aristro Grid Lamp Products, Inc.  
65 Harbor Road  
Fort Washington, NY 11050

CooperVision Diagnostics  
7356 Trade Street  
San Diego, CA 92121

Mentor O & O Inc.  
South Shore Park  
Hingham, MA 02043

Nicolet Biomedical Division  
5225-4 Verona Road  
P. O. Box 4287  
Madison, WI 53711-0287

Reichert Scientific Instruments  
P. O. Box 123  
Buffalo, NY 14240

### INITIAL DISTRIBUTION

Commander  
US Army Natick Research and  
Development Center  
ATTN: Documents Librarian  
Natick, MA 01760

Commander  
US Army Research Institute of  
Environmental Medicine  
Natick, MA 01760

Naval Submarine Medical Research  
Laboratory  
Medical Library, Naval Sub Base  
Box 900  
Groton, CT 05340

US Army Avionics Research and  
Development Activity  
ATTN: SAVAA-P-TP  
Fort Monmouth, NJ 07703-5401

Commander/Director  
US Army Combat Surveillance and  
Target Acquisition Laboratory  
ATTN: DELCS-D  
Fort Monmouth, NJ 07703-5304

US Army Research and Development  
Support Activity  
Fort Monmouth, NJ 07703

ATTN: Dr.  
US Army Research and Development  
Support Activity  
Fort Monmouth, NJ 07703

US Army Research and Development  
Support Activity  
Fort Monmouth, NJ 07703

US Army Research and Development  
Support Activity  
Fort Monmouth, NJ 07703

Commander  
Man-Machine Integration System  
Code 602  
Naval Air Development Center  
Warminster, PA 18974

Naval Air Development Center  
Technical Information Division  
Technical Support Detachment  
Warminster, PA 18974

Commander  
Naval Air Development Center  
ATTN: Code 6021 (Mr. Brindle)  
Warminster, PA 18974

Dr. E. Hendler  
Human Factors Applications, Inc.  
295 West Street Road  
Warminster, PA 18974

Commanding Officer  
Naval Medical Research and  
Development Command  
National Naval Medical Center  
Bethesda, MD 20014

Under Secretary of Defense for  
Research and Engineering  
ATTN: Military Assistant for  
Medical and Life Sciences  
Washington, DC 20301

Director  
Army Audiology and Speech Center  
Walter Reed Army Medical Center  
Washington, DC 20307-5001

COL Franklin H. Top, Jr., MD  
Walter Reed Army Institute  
of Research  
Washington, DC 20307-5100

Commander  
US Army Institute of Dental Research  
Walter Reed Army Medical Center  
Washington, DC 20307-5300



HQ DA (DASG-PSP-O)  
Washington, DC 20310

Naval Air Systems Command  
Technical Library Air 950D  
Rm 278, Jefferson Plaza II  
Department of the Navy  
Washington, DC 20361

Naval Research Laboratory Library  
Code 1433  
Washington, DC 20375

Naval Research Laboratory Library  
Shock & Vibration Information Center  
Code 5804  
Washington, DC 20375

Harry Diamond Laboratories  
ATTN: Tech Information Branch  
2800 Powder Mill Road  
Adelphi, MD 20783-1197

Director  
US Army Human Engineering Laboratory  
ATTN: Technical Library  
Aberdeen Proving Ground, MD  
21005-5001

US Army Materiel Systems  
Analysis Agency  
ATTN: Reports Processing  
Aberdeen Proving Ground, MD  
21005-5017

Commander  
US Army Test & Evaluation Command  
ATTN: AMSTE-AD-H  
Aberdeen Proving Ground, MD  
21005-5055

US Army Ordnance Center  
& School Library  
Bldg 3071  
Aberdeen Proving Ground, MD  
21005-5201

Director  
US Army Ballistic Research Laboratory  
ATTN: DRXBR-OD-ST Tech Reports  
Aberdeen Proving Ground, MD  
21005-5066

US Army Environmental Hygiene  
Agency Library  
Bldg E2100  
Aberdeen Proving Ground, MD 21010

Commander  
US Army Medical Research Institute  
of Chemical Defense  
ATTN: SGRD-UV-AO  
Aberdeen Proving Ground, MD  
21010-5425

Technical Library  
Chemical Research & Development Center  
Aberdeen Proving Ground, MD  
21010-5423

Commander  
US Army Medical Research  
& Development Command  
ATTN: SGRD-RMS (Mrs. Madigan)  
Fort Detrick, Frederick, MD  
21701-5012

Commander  
US Army Medical Research Institute  
of Infectious Diseases  
Fort Detrick, Frederick, MD 21701

Commander  
US Army Medical Bioengineering  
Research & Development Laboratory  
ATTN: SGRD-UBZ-I  
Fort Detrick, Frederick, MD 21701

Dr. R. Newburgh  
Director, Biological Sciences Division  
Office of Naval Research  
600 North Quincy Street  
Arlington, VA 22217

Defense Technical Information Center  
Cameron Station  
Alexandria, VA 22314

Commander  
US Army Materiel Command  
ATTN: AMCDE-S (CPT Broadwater)  
5001 Eisenhower Avenue  
Alexandria, VA 22333

US Army Foreign Science and  
Technology Center  
ATTN: MTZ  
220 7th Street, NE  
Charlottesville, VA 22903-0096

Commandant  
US Army Aviation Logistics School  
ATTN: ATSQ-TDN  
Fort Eustis, VA 23604

Director, Applied Technology Lab  
USARTL-AVSCOM  
ATTN: Library, Bldg 401  
Fort Eustis, VA 23604

US Army Training and  
Doctrine Command  
ATTN: ATCD-ZX  
Fort Monroe, VA 23651

US Army Training and  
Doctrine Command  
ATTN: Surgeon  
Fort Monroe, VA 23651-5000

Structures Laboratory Library  
USARTL-AVSCOM  
NASA Langley Research Center  
Mail Stop 266  
Hampton, VA 23665

Aviation Medicine Clinic  
TMC #22, SAAF  
Fort Bragg, BC 28305

Naval Aerospace Medical  
Institute Library  
Bldg 1953, Code 102  
Pensacola, FL 32508

US Air Force Armament Development  
and Test Center  
Eglin Air Force Base, FL 32542

Command Surgeon  
US Central Command  
MacDill AFB, FL 33608

US Army Missile Command  
Redstone Scientific Information Center  
ATTN: Documents Section  
Redstone Arsenal, AL 35898-5241

Air University Library  
AUL LSEN  
Maxwell AFB, AL 36112

Commander  
US Army Aeromedical Center  
Fort Rucker, AL 36362

Commander  
US Army Aviation Center & Fort Rucker  
ATTN: ATZQ-CDR  
Fort Rucker, AL 36362

Directorate of Combat Developments  
Bldg 507  
Fort Rucker, AL 36362

Directorate of Training Development  
Bldg 502  
Fort Rucker, AL 36362

Chief  
Army Research Institute Field Unit  
Fort Rucker, AL 36362

Chief  
Human Engineering Labs Field Unit  
Fort Rucker, AL 36362

Commander  
US Army Safety Center  
Fort Rucker, AL 36362

Commander  
US Army Aviation Center & Fort Rucker  
ATTN: ATZQ-T-ATL  
Fort Rucker, AL 36362

US Army Aircraft Development  
Test Activity  
ATTN: STEBG-MP-QA  
Cairns AAF, Ft Rucker, AL 36362

President  
US Army Aviation Board  
Cairns AAF, Ft Rucker, AL 36362

US Army Research & Technology  
Laboratories (AVSCOM)  
Propulsion Laboratory MS 302-2  
NASA Lewis Research Center  
Cleveland, OH 44135

AFAMRL/HEX  
Wright-Patterson AFB, OH 45433

US Air Force Institute of Technology  
(AFIT/LDEE)  
Bldg 640, Area B  
Wright-Patterson AFB, OH 45433

University of Michigan  
NASA Center of Excellence  
in Man-Systems Research  
ATTN: R.G. Snyder, Director  
Ann Arbor, MI 48109

Henry L. Taylor  
Director, Institute of Aviation  
Univ of Illinois - Willard Airport  
Savoy, IL 61874

John A. Dellinger, MS, ATP  
Univ of Illinois - Willard Airport  
Savoy, IL 61874

Commander  
US Army Aviation Systems Command  
ATTN: DRSAB-WS  
4300 Goodfellow Blvd  
St Louis, MO 63120-1798

Project Officer  
Aviation Life Support Equipment  
ATTN: AMCPO-ALSE  
4300 Goodfellow Blvd  
St Louis, MO 63120-1798

Commander  
US Army Aviation Systems Command  
ATTN: SGRD-UAX-AL (MAJ Lacy)  
Bldg 105, 4300 Goodfellow Blvd  
St Louis, MO 63120

Commander  
US Army Aviation Systems Command  
ATTN: DRSAB-ED  
4300 Goodfellow Blvd  
St Louis, MO 63120

US Army Aviation Systems Command  
Library & Info Center Branch  
ATTN: DRSAB-DIL  
4300 Goodfellow Blvd  
St Louis, MO 63120

Commanding Officer  
Naval Biodynamics Laboratory  
P.O. Box 24907  
New Orleans, LA 70189

Federal Aviation Administration  
Civil Aeromedical Institute  
CAMI Library AAC 64D1  
P.O. Box 25082  
Oklahoma City, OK 73125

US Army Field Artillery School  
ATTN: Library  
Snow Hall, Room 14  
Fort Sill, OK 73503

Commander  
US Army Academy of Health Sciences  
ATTN: Library  
Fort Sam Houston, TX 78234

Commander  
US Army Health Services Command  
ATTN: HSOP-SO  
Fort Sam Houston, TX 78234-6000

Commander  
US Army Institute of Surgical Research  
ATTN: SGRD-USM (Jan Duke)  
Fort Sam Houston, TX 78234-6200

Director of Professional Services  
AFMSC/GSP  
Brooks Air Force Base, TX 78235

US Air Force School  
of Aerospace Medicine  
Strughold Aeromedical Library  
Documents Section, USAFSAM/TSK-4  
Brooks Air Force Base, TX 78235

US Army Dugway Proving Ground  
Technical Library  
Bldg 5330  
Dugway, UT 84022

Dr. Diane Damos  
Department of Human Factors  
ISSM, USC  
Los Angeles, CA 90089-0021

US Army Yuma Proving Ground  
Technical Library  
Yuma, AZ 85364

US Army White Sands Missile Range  
Technical Library Division  
White Sands Missile Range, NM 88002

US Air Force Flight Test Center  
Technical Library, Stop 238  
Edwards Air Force Base, CA 93523

US Army Aviation Engineering  
Flight Activity  
ATTN: SAVTE-M (Tech Lib) Stop 217  
Edwards AFB, CA 93523-5000

Commander  
Code 3431  
Naval Weapons Center  
China Lake, CA 93555

US Army Combat Developments  
Experimental Center  
Technical Information Center  
Bldg 2925  
Fort Ord, CA 93941-5000

Aeromechanics Laboratory  
US Army Research  
& Technical Laboratories  
Ames Research Center, M/S 215-1  
Moffett Field, CA 94035

Commander  
Letterman Army Institute of Research  
ATTN: Medical Research Library  
Presidio of San Francisco, CA 94129

Sixth US Army  
ATTN: SMA  
Presidio of San Francisco, CA 94129

Director  
Naval Biosciences Laboratory  
Naval Supply Center, Bldg 844  
Oakland, CA 94625

Col G. Stebbing  
USDAO-AMLO, US Embassy  
Box 36  
FPO New York 09510

Staff Officer, Aerospace Medicine  
RAF Staff, British Embassy  
3100 Massachusetts Avenue, NW  
Washington, DC 20008

Canadian Society of Aviation Medicine  
c/o Academy of Medicine, Toronto  
ATTN: Ms. Carmen King  
288 Bloor Street West  
Toronto, Ontario M5S 1V8

Canadian Airline Pilot's Association  
MAJ J. Soutendam (Retired)  
1300 Steeles Avenue East  
Brampton, Ontario, L6T 1A2

Canadian Forces Medical Liaison Officer  
Canadian Defence Liaison Staff  
2450 Massachusetts Avenue, NW  
Washington, DC 20008

Commanding Officer  
404 Squadron CFB Greenwood  
Greenwood, Nova Scotia BOP 1N0

Officer Commanding  
School of Operational  
& Aerospace Medicine  
DCIEM, P.O. Box 2000  
1133 Sheppard Avenue West  
Downsview, Ontario M3M 3B9

National Defence Headquarters  
101 Colonel By Drive  
ATTN: DPM  
Ottawa, Ontario K1A 0K2

Commanding Officer  
Headquarters, RAAF Base  
POINT COOK VIC 3029  
Australia

Canadian Army Liaison Office  
Bldg 602  
Fort Rucker, AL 36362

Netherlands Army Liaison Office  
Bldg 602  
Fort Rucker, AL 36362

German Army Liaison Office  
Bldg 602  
Fort Rucker, AL 36362

British Army Liaison Office  
Bldg 602  
Fort Rucker, AL 36362

French Army Liaison Office  
Bldg 602  
Fort Rucker, AL 36362

END

DTIC

6-86